

**With Organic Soil Tolerance for Veterinary Clinics, Animal Hospitals, Pet Shops, Kennels, Animal Care Facilities, Tack Shops, Animal Life Science Laboratories, Breeding and Grooming Establishments and Farms.**

**KENNEL MASTER** is designed for use in veterinary clinics, animal hospitals, animal laboratories, animal life science laboratories, animal care facilities, kennels, tack shops, pet shops, zoos, pet animal quarters, stables, farms and pet grooming/breeding establishments. When used as directed, this formulation is specifically designed to disinfect, deodorize and clean inanimate, hard, non-porous surfaces: walls, floors, sink tops, furniture, operating tables, kennel runs, cages, and feeding and watering equipment. In addition, **KENNEL MASTER** will deodorize those areas that are generally hard to keep fresh smelling such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

**\*\*Kills Pseudomonas aeruginosa, Escherichia coli, Klebsiella pneumoniae, Salmonella schottmuelleri, Salmonella choleraesuis, Streptococcus faecalis, Shigella dysenteriae, Enterobacter aerogenes, Staphylococcus aureus, Bordetella bronchiseptica, Canine Distemper virus, Pseudorabies virus, Vaccinia virus, Influenza A2, Herpes Simplex Type 1, Adenovirus Type 2, Bovine Viral Diarrhea Virus (BVDV), Norovirus (Norwalk Virus), Feline Calicivirus (FCV), Canine Parvovirus (CPV), Canine Adenovirus, Canine Coronavirus, Canine Parainfluenza Virus, Feline Coronavirus, Feline Rhinotracheitis Virus, Feline Picornavirus, Feline Panleukopenia and Porcine Parvovirus.**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DISINFECTION** - For disinfecting, add 1 ounce of **KENNEL MASTER** to one gallon of water. Apply this solution with a mop, cloth, sponge, hand pump trigger sprayer, or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow surfaces to remain wet for **10 minutes** and then let air dry.

**BACTERICIDAL ACTIVITY** - When used as directed, **KENNEL MASTER** demonstrates effective disinfectant activity against *Pseudomonas aeruginosa*, *Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella schottmuelleri*, *Salmonella choleraesuis*, *Streptococcus faecalis*, *Shigella dysenteriae*, *Enterobacter aerogenes* and *Staphylococcus aureus*. When used as directed, this formulation is an effective disinfectant against *Pseudomonas aeruginosa* in animal hospitals and laboratories. At 4 ounces per gallon or equivalent dilution rate (2812 ppm (mg/L) active quat), **KENNEL MASTER** is effective against *Bordetella bronchiseptica* (causative agent of bacterial Kennel Cough) on hard, non-porous environmental surfaces.

**\*VIRUCIDAL ACTIVITY** - When used at the 1 ounce to one gallon or equivalent dilution rate (703 ppm (mg/L) active quat), **KENNEL MASTER** exhibits effective virucidal activity on inanimate, hard, non-porous surfaces against Canine Distemper virus, Pseudorabies virus, Vaccinia virus, Influenza A2, Herpes Simplex Type 1, Adenovirus Type 2, Bovine Viral Diarrhea Virus (BVDV), Norovirus (Norwalk Virus), and Feline Calicivirus (FCV). At 4 ounces per gallon or equivalent dilution rate (2812 ppm (mg/L) active quat), **KENNEL MASTER** is effective against the following viruses on hard, non-porous environmental surfaces: Canine Parvovirus (CPV), Canine Adenovirus (causative agent of Infectious Canine Hepatitis), Canine Coronavirus (causative agent of Canine Coronavirus Disease, Canine Gastroenteritis), Canine Parainfluenza Virus (causative agent of viral Kennel Cough), Feline Panleukopenia (Feline Parvovirus), Feline Rhinotracheitis (causative agent of Feline Herpes virus Infection), Feline Coronavirus (causative agent of Feline Infectious Peritonitis), Feline Picornavirus and Porcine Parvovirus. **Canine Parvovirus (CPV) Activity:** At 4 ounces per gallon of water or equivalent dilution, **KENNEL MASTER** inactivates Canine Parvovirus (CPV) on hard, non-porous, environmental surfaces in the presence of 5% blood serum with a **10 minute** contact time.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

**MILDEWSTAT** - To control mold and mildew on pre-cleaned hard, non-porous surfaces, add 1 ounce to one gallon of water (or equivalent dilution). Apply use-solution with a cloth, mop, sponge, hand pump trigger sprayer or low pressure coarse sprayer making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.



WECHEM, INC.  
5734 Susitna Dr.  
Harahan, LA 70123  
Local: (504) 733-1152  
USA TOLL FREE: 1-800-426-0512  
FAX: (504) 733-2218 \* www.wechem.com

**KENNEL MASTER™**

**Detergent; Disinfectant; Deodorant;  
\*Virucide; Mildewstat  
Kills Germs\*\***

**Kills Pandemic 2009 H1N1 Influenza A Virus**

**ACTIVE INGREDIENTS**

<b>n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)</b>	
<b>dimethyl benzyl ammonium chlorides .....</b>	<b>4.5%</b>
<b>n-Alkyl (68% C12, 32% C14)</b>	
<b>dimethyl ethyl benzyl ammonium chlorides.....</b>	<b>4.5%</b>
<b>INERT INGREDIENTS .....</b>	<b>91.0%</b>
<b>TOTAL .....</b>	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

**See side panel for additional precautionary statements.**

**EPA REG. NO. 1839-100-34370 EPA EST. NO. 34370-LA-001**

**NET CONTENTS: 1 U.S. GALLON**

**BATCH**

**07/10**

**DEODORIZATION** - To deodorize, add 1 ounce of **KENNEL MASTER** per gallon of water or equivalent dilution of water (703 ppm (mg/L) active quat). Excess material should be wiped up or allowed to air dry. **For pet odors:** Use on carpets, floors, walls, tile, cages, pet beds, litter boxes, mats, floor coverings, area rugs or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area, then follow directions for DEODORIZATION.

**VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS / BREEDING AND GROOMING ESTABLISHMENTS / TACK SHOPS / ANIMAL LIFE SCIENCE LABORATORIES DISINFECTION & \*VIRUCIDAL ACTIVITY DIRECTIONS** - For cleaning and disinfecting the following hard, non-porous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feed from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 4 ounces of **KENNEL MASTER** per gallon of water (or equivalent dilution) for a period of **10 minutes**. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other equipment that dispenses food or water with soap or detergent, and rinse with potable water before reuse.

**FARM PREMISE DISINFECTION DIRECTIONS (For Use in Equine, Dairy and Hog Farms) -**

- Do not use in milking stalls, milking parlors or milk houses.
- 1. Remove all animals and feed from premises, vehicles and enclosures.
- 2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals.
- 3. Empty all troughs, racks and other feeding and watering appliances.
- 4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
- 5. Saturate all surfaces with a use-solution of 4 ounces of **KENNEL MASTER** per gallon of water (or equivalent dilution) for a period of **10 minutes**.
- 6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure.
- 7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried.
- 8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE.** Causes irreversible eye damage and skin burns. Do not get in eyes on skin or on clothing. May be fatal if absorbed through skin. Harmful if swallowed. Wear goggles or face shield, rubber gloves, and protective clothing. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash before reuse.

**STORAGE AND DISPOSAL**

**DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.**

**PESTICIDE STORAGE** - Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in an area inaccessible to children.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL** - Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying. Triple rinse as follows: Fill container 1/4 full with water and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for **10 seconds** after the flow begins to drip. Repeat procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.