

**Pro-Line** is a phosphate-free, pH neutral formulation designed to provide effective cleaning, deodorizing, and disinfection for hospitals, nursing homes, schools, transportation terminals, office buildings, manufacturing facilities, lodging establishments, hotels, retail businesses, and athletic/recreation facilities, where housekeeping is of prime importance in controlling cross-contamination from treated surfaces. **Pro-Line** is a one step neutral disinfectant that is effective against a broad spectrum of bacteria, is virucidal\*(including HIV-1, HIV-2, HBV and HCV) and inhibits the growth of mold and mildew and their odors when used as directed. **Pro-Line**, when used as directed, is formulated to disinfect the following hard non-porous inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, sealed fiberglass, sealed granite, sealed marble, chrome, vinyl, plastic surfaces, vanity tops, shower stalls, bathtubs, and cabinets.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical and semi-critical medical device surfaces.

**DISINFECTION** - To disinfect inanimate, hard non-porous surfaces add 4 ounces of **Pro-Line** per gallon of water. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain wet for **10 minutes**, then remove excess liquid. For sprayer applications, spray 6-8 inches from surface, rub with brush, sponge or cloth. Do not breathe spray mist. For heavily soiled areas, a pre-cleaning step is required.

**BACTERICIDAL STABILITY OF USE-SOLUTION** - Tests confirm that **Pro-Line** when diluted at 4 ounces/gallon in hard water up to 400 ppm (as CaCO<sub>3</sub>) remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella enterica* for up to 1 month when stored in a sealed container such as a spray bottle at room temperature. If **Pro-Line** becomes visibly dirty or contaminated, the use solution must be discarded and fresh use solution prepared. Always use clean properly labeled containers when diluting **Pro-Line**. Bactericidal stability of the use-solution does not apply to open containers such as buckets or pails.

**To disinfect toilet bowls** - Remove gross filth or soils from surfaces with bowl brush. Add 4 ounces of **Pro-Line** to the bowl water. Brush or swab the bowl completely using a scrub brush or toilet mop making sure to get under the rim. Let stand for **10 minutes** and flush.

**BACTERICIDAL ACTIVITY** - At the 4 ounce per gallon dilution, this product demonstrates effective disinfectant activity against the organisms: *Pseudomonas aeruginosa* PRD-10, *Salmonella* (choleraesuis) enterica, *Staphylococcus aureus*, *Staphylococcus aureus* (clinical isolate), *Bordetella bronchiseptica*, *Corynebacterium ammoniagenes*, *Enterobacter aerogenes*, *Enterobacter cloacae*, *Enterobacter cloacae* (clinical isolate), *Enterococcus faecalis*, *Enterococcus faecalis* (clinical isolate), *Escherichia coli*, *Escherichia coli* (clinical isolate), *Fusobacterium necrophorum*, *Klebsiella pneumoniae* subsp. pneumoniae, *Lactobacillus casei* subsp. ramosus, *Listeria monocytogenes*, *Pasteurella multocida*, *Proteus vulgaris*, *Proteus mirabilis* ATCC 9921, *Proteus mirabilis* ATCC 25933, *Salmonella* (paratyphi B) enterica, *Salmonella* (typhi) enterica, *Salmonella* (typhimurium) enterica, *Salmonella* (pullorum) enterica, *Serratia marcescens*, *Shigella sonnei*, *Shigella flexneri* Type 2b, *Shigella dysenteriae*, *Staphylococcus aureus* subsp. aureus, *Staphylococcus epidermidis*, *Staphylococcus epidermidis* (clinical isolate), *Streptococcus pyogenes* (Clinical - Flesh Eating Strain BIRD M3), *Streptococcus pyogenes* Group A, *Xanthomonas maltophilia* (clinical isolate), *Vancomycin* resistant *Enterococcus faecalis* (VRE), *Methicillin* resistant *Staphylococcus aureus* (MRSA), *Vancomycin* intermediate resistant *Staphylococcus aureus* (VISA), *Community Associated Methicillin* resistant *Staphylococcus aureus* (CA-MRSA) NRS 123 Genotype USA400, *Community Associated Methicillin* resistant *Staphylococcus aureus* (CA-MRSA) NRS 384 Genotype USA300, *Salmonella* (enteritidis) enterica, *Ampicillin* resistant *Acinetobacter baumannii*, *Gefazolin* resistant *Acinetobacter baumannii*, *Ceftazidime* resistant *Acinetobacter baumannii*, *Ceftaroxime* resistant *Acinetobacter baumannii*, *Gentamicin* resistant *Acinetobacter baumannii*, *Tobramycin* resistant *Acinetobacter baumannii*, *Ciprofloxacin* resistant *Acinetobacter baumannii*, *Levofloxacin* resistant *Acinetobacter baumannii* and *Bactrim* resistant *Acinetobacter baumannii*.

**MILDEWSTAT** - To control mold and mildew (*Aspergillus niger*) and the odors they cause on pre-cleaned, hard, non-porous inanimate surfaces add 4 ounces of this product per gallon of water. Apply solution with a cloth, mop, sponge or hand pump trigger sprayer making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

**FUNGICIDAL ACTIVITY** - At the 4 ounce per gallon dilution, **Pro-Line** is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* (Athlete's Foot fungus) (a cause of ringworm) and *Candida albicans*. Apply solution with a cloth, sponge or hand pump trigger sprayer to hard, non-porous surfaces such as shower stalls, locker rooms, exercise floors or other clean, hard, non-porous surfaces commonly contacted by bare feet. Allow the surface to remain wet for **10 minutes**, then remove excess liquid. Diluted product should be applied daily or more frequently with heavy facility use.

**\*VIRUCIDAL ACTIVITY** - This product, when used on environmental, inanimate, hard, non-porous surfaces, exhibits effective virucidal activity against HIV-1, HIV-2, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Type 1 (causative agent of fever blisters), Herpes Simplex Type 2 (genital), Influenza A2/Hong Kong, Vaccinia, Rotavirus, Human Coronavirus (ATCC VR-740, Strain 229E), SARS associated Coronavirus, Bovine Viral Diarrhea Virus (BVDV), Pseudorabies, Bovine Rhinotracheitis, Feline Leukemia, Feline Picornavirus, Avian Influenza A Virus H9N2, Avian Influenza A Virus H3N2, Canine Distemper Virus, Rabies Virus, Paramyxovirus (Mumps) and Porcine Respiratory & Reproductive Syndrome Virus (PRRSV). Add 4 ounces of this product per gallon of water. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a cloth, mop, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow the surface to remain wet for **10 minutes**, then remove excess liquid. **KILLS HIV-1, HIV-2, HBV, and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood

**WECHEM®** *Engineered Chemistries*  
**SUPERIOR SOLUTIONS**

# PRO-LINE™

Cleaner; Detergent; Disinfectant; Deodorizer; \*Virucide;  
Fungicide (against pathogenic fungi);  
Mildewstat (on hard, non-porous, inanimate surfaces)

## ACTIVE INGREDIENTS

Octyl decyl dimethyl ammonium chloride ..... 0.814%

Dioctyl dimethyl ammonium chloride ..... 0.407%

Didecyl dimethyl ammonium chloride ..... 0.407%

Alkyl (50% C14, 40% C12, 10% C16)

dimethyl benzyl ammonium chloride ..... 1.085%

INERT INGREDIENTS ..... 97.287%

TOTAL ..... 100.000%

KEEP OUT OF REACH OF CHILDREN  
**DANGER**

## FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call poison control center or doctor for treatment advice.

**If swallowed:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

See side panel for additional precautionary statements.

EPA REG. NO. 1839-168-34370

WECHEM, INC.

5734 Susitna Dr. Harahan, LA 70123

Local: (504) 733-1152 \* USA TOLL FREE: 1-800-426-0512

FAX: (504) 733-2218 \* www.wechem.com

EPA EST. NO. 34370-LA-001

NET CONTENTS: 55 U.S. GALLONS

BATCH

6/11

of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/object likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 or Type 2 (HIV-1 or HIV-2) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HIV-2, HBV, and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** Allow surface to remain wet for **10 minutes**.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

**GENERAL CLEANING:** Apply this product to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer. This product can be diluted at 4 ounces per gallon, or applied at full strength. Thoroughly wet soiled surface and scrub wet surface.

**DEODORIZATION:** To deodorize, apply this product as indicated under the heading DISINFECTION.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE AND \*VIRUCIDE AGAINST THE LISTED ORGANISMS IN WATER UP TO 400 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM). THIS PRODUCT IS AN EFFECTIVE FUNGICIDE AGAINST THE LISTED FUNGI IN WATER UP TO 200 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

(Not for use in California)

**WATER AND SMOKE DAMAGE RESTORATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI FOR INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE.**

**Pro-Line** is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim and frame lumber, tackless strip and paneling. Mix 4 ounces of this product per gallon of water, saturate affected materials with enough use-solution to remain wet for at least **10 minutes**. Use proper ventilation.

**Sewer backup & river flooding:** During mitigation procedures, dilute 4 to 8 ounces of this product per gallon of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affected areas with a sprayer using a coarse spray tip, before and after cleaning and extraction.

**Carpets, carpet cushions and other porous materials such as sub floors, drywall, trim and frame lumber, tackless strip and paneling:** For water damage from a clean water source, extract excess water. Test hidden area for colorfastness. Mix 4 to 8 ounces of this product per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a spray using a coarse spray tip, to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for **10 minutes**. Follow with a thorough extraction. Dry rapidly and thoroughly.

**NOTE:** This product must not be mixed with other cleaning products.

## PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE.** Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. May be fatal if absorbed through the skin. Harmful if swallowed. Wear goggles or face shield, rubber gloves, and protective clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

## STORAGE AND DISPOSAL

**DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.**

**PESTICIDE STORAGE** - Store in a dry place no lower in temperature than 50°F or higher than 120°F. Store in an area inaccessible to children.

**CONTAINER DISPOSAL** - Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container promptly after emptying. Triple rinse as follows: Fill container 1/4 full with water. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Follow Pesticide Disposal instructions for rinse disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning. If not available, puncture and dispose in a sanitary landfill.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

